

# Some Applications of 2-D Systems of Differential Equations

Xu-Yan Chen

Examples:

- ▶ Salt in Tanks (a linear system)
- ▶ Electric Circuits (a linear system)
- ▶ Population Model - Competing Species (a nonlinear system)

# Linear Model of Population Dynamics

Malthus (1798)

Diff equation for the population  $P(t)$  at time  $t$ :

$$P' = rP,$$

where constant  $r$  is the net per capita growth rate:

$$r = b - d = \text{per capita birth rate} - \text{per capita death rate}.$$

**Solutions:**  $P(t) = P(0)e^{rt}$

---

When  $r > 0$ ,


## Phase portrait



Equilibrium  $P = 0$  is unstable.

Positive solutions  $P(t)$  grow exponentially to  $\infty$  as  $t \rightarrow \infty$ .

## Solution Graphs $P(t)$ vs $t$



# Logistic Model of Population Dynamics

Verhulst (1838)

$$P' = rP \left( 1 - \frac{P}{K} \right)$$


where  $r$  is the net per capita growth rate when  $P \approx 0$ ,  
 $K$  is the *carrying capacity*.

**Solution Formula:** 
$$P(t) = \frac{KP(0)}{P(0) + [K - P(0)]e^{-rt}}$$

---

Example.  $P' = 6P(1 - P/2)$   $(r = 6, K = 2)$


## Phase portrait



Equilibrium  $P = 0$  is unstable.

Equilibrium  $P = K$  is asymp stable.

All positive solutions  $P(t)$  converge to  $K$  as  $t \rightarrow \infty$ .



# Logistic Dynamics of Two Species


If no interactions: 
$$\begin{cases} x'_1 = x_1(6 - 3x_1) \\ x'_2 = x_2(3 - 2x_2) \end{cases}$$

---

# Logistic Dynamics of Two Species

If no interactions:  $\begin{cases} x'_1 = x_1(6 - 3x_1) \\ x'_2 = x_2(3 - 2x_2) \end{cases} \implies \begin{aligned} \lim_{t \rightarrow \infty} x_1(t) &= 2 \\ \lim_{t \rightarrow \infty} x_2(t) &= 3/2 \end{aligned}$


---



# Logistic Dynamics of Two Species

If no interactions:  $\begin{cases} x'_1 = x_1(6 - 3x_1) \\ x'_2 = x_2(3 - 2x_2) \end{cases} \implies \begin{aligned} \lim_{t \rightarrow \infty} x_1(t) &= 2 \\ \lim_{t \rightarrow \infty} x_2(t) &= 3/2 \end{aligned}$

---




# Logistic Dynamics of Two Species

If no interactions:  $\begin{cases} x'_1 = x_1(6 - 3x_1) \\ x'_2 = x_2(3 - 2x_2) \end{cases} \implies \begin{aligned} \lim_{t \rightarrow \infty} x_1(t) &= 2 \\ \lim_{t \rightarrow \infty} x_2(t) &= 3/2 \end{aligned}$

---

Nonlinear Dynamics





# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

**With Competition:**

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

**With Competition:**

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

**Basic Questions:**

# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

**With Competition:**

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

**Basic Questions:**

- Find equilibria. (i.e., time independent solutions)

# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

## With Competition:

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

## Basic Questions:

- Find equilibria. (i.e., time independent solutions)
- Construct a linear approximating system near each equilibrium.  
(use the Jacobian matrix, that is, partial derivatives)

# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

## With Competition:

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

## Basic Questions:

- ▶ Find equilibria. (i.e., time independent solutions)
- ▶ Construct a linear approximating system near each equilibrium.  
(use the Jacobian matrix, that is, partial derivatives)
- ▶ Study the linear approximating dynamics near the equilibrium.  
(use eigenvalues & eigenvectors)

# Example 3. Logistic Growth & Competition

Lotka (1925), Volterra (1926), Gause (1934), . . . . .

## With Competition:

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - 2x_2 - x_1) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

## Basic Questions:

- ▶ Find equilibria. (i.e., time independent solutions)
- ▶ Construct a linear approximating system near each equilibrium.  
(use the Jacobian matrix, that is, partial derivatives)
- ▶ Study the linear approximating dynamics near the equilibrium.  
(use eigenvalues & eigenvectors)
- ▶ Determine the nonlinear dynamics near the equilibrium.  
(if eigenvalues are  $\neq 0$  & are not purely imaginary, Yes We Can!)

## Example 3. (Continued. Find equilibria.)

**Competing Species:**

$$\begin{cases} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{cases}$$

$$\left| \begin{aligned} f_1(x_1, x_2) &= x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) &= x_2(3 - x_1 - 2x_2) \end{aligned} \right.$$

## Example 3. (Continued. Find equilibria.)

**Competing Species:**

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

**Equilibria:**

$$\left\{ \begin{array}{l} x_1(6 - 3x_1 - 2x_2) = 0 \\ x_2(3 - x_1 - 2x_2) = 0 \end{array} \right.$$



## Example 3. (Continued. Find equilibria.)

**Competing Species:**

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

**Equilibria:**

$$\left\{ \begin{array}{l} x_1(6 - 3x_1 - 2x_2) = 0 \\ x_2(3 - x_1 - 2x_2) = 0 \end{array} \right. \implies \text{Separate to four combinations}$$

$$\left\{ \begin{array}{l} x_1 = 0 \\ x_2 = 0 \end{array} \right. \quad \left\{ \begin{array}{l} 6 - 3x_1 - 2x_2 = 0 \\ x_2 = 0 \end{array} \right.$$

$$\left\{ \begin{array}{l} x_1 = 0 \\ 3 - x_1 - 2x_2 = 0 \end{array} \right. \quad \left\{ \begin{array}{l} 6 - 3x_1 - 2x_2 = 0 \\ 3 - x_1 - 2x_2 = 0 \end{array} \right.$$

## Example 3. (Continued. Find equilibria.)

**Competing Species:**

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

**Equilibria:**

$$\left\{ \begin{array}{l} x_1(6 - 3x_1 - 2x_2) = 0 \\ x_2(3 - x_1 - 2x_2) = 0 \end{array} \right. \implies \text{Separate to four combinations}$$

$$\left\{ \begin{array}{l} x_1 = 0 \\ x_2 = 0 \end{array} \right. \quad \left\{ \begin{array}{l} 6 - 3x_1 - 2x_2 = 0 \\ x_2 = 0 \end{array} \right.$$

$$\left\{ \begin{array}{l} x_1 = 0 \\ 3 - x_1 - 2x_2 = 0 \end{array} \right. \quad \left\{ \begin{array}{l} 6 - 3x_1 - 2x_2 = 0 \\ 3 - x_1 - 2x_2 = 0 \end{array} \right.$$

Four equilibria:

$$(x_1, x_2) = (0, 0), \quad (2, 0), \quad (0, \tfrac{3}{2}), \quad (\tfrac{3}{2}, \tfrac{3}{4}).$$

## Example 3. (continued. Linear Approximation)

Competing Species:

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

## Example 3. (continued. Linear Approximation)

Competing Species:

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

Linear Approximating System near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

## Example 3. (continued. Linear Approximation)

Competing Species:

$$\left\{ \begin{array}{l} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

Linear Approximating System near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

- Prepare the Jacobian matrix:

$$J = \begin{bmatrix} \partial f_1 / \partial x_1 & \partial f_1 / \partial x_2 \\ \partial f_2 / \partial x_1 & \partial f_2 / \partial x_2 \end{bmatrix} = \begin{bmatrix} 6 - 6x_1 - 2x_2 & -2x_1 \\ -x_2 & 3 - x_1 - 4x_2 \end{bmatrix}$$

## Example 3. (continued. Linear Approximation)

Competing Species:

$$\left\{ \begin{array}{l} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

Linear Approximating System near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

- Prepare the Jacobian matrix:

$$J = \begin{bmatrix} \partial f_1 / \partial x_1 & \partial f_1 / \partial x_2 \\ \partial f_2 / \partial x_1 & \partial f_2 / \partial x_2 \end{bmatrix} = \begin{bmatrix} 6 - 6x_1 - 2x_2 & -2x_1 \\ -x_2 & 3 - x_1 - 4x_2 \end{bmatrix}$$

- Evaluate  $J$  at equilibrium  $(x_1, x_2) = (\frac{3}{2}, \frac{3}{4})$ :

$$J = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix}$$

## Example 3. (continued. Linear Approximation)

Competing Species:

$$\left\{ \begin{array}{l} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - x_1 - 2x_2) \end{array} \right. \quad \left| \quad \begin{array}{l} f_1(x_1, x_2) = x_1(6 - 3x_1 - 2x_2) \\ f_2(x_1, x_2) = x_2(3 - x_1 - 2x_2) \end{array} \right.$$

Linear Approximating System near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

- Prepare the Jacobian matrix:

$$J = \begin{bmatrix} \partial f_1 / \partial x_1 & \partial f_1 / \partial x_2 \\ \partial f_2 / \partial x_1 & \partial f_2 / \partial x_2 \end{bmatrix} = \begin{bmatrix} 6 - 6x_1 - 2x_2 & -2x_1 \\ -x_2 & 3 - x_1 - 4x_2 \end{bmatrix}$$

- Evaluate  $J$  at equilibrium  $(x_1, x_2) = (\frac{3}{2}, \frac{3}{4})$ :

$$J = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix}$$

- The Linear Approximating System near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix} \begin{bmatrix} x_1 - \frac{3}{2} \\ x_2 - \frac{3}{4} \end{bmatrix}$$

## Example 3. Linear dynamics near $(0, 0)$

The Linear Approximating  
System near equilibrium  $(0, 0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$



## Example 3. Linear dynamics near $(0, 0)$

The Linear Approximating  
System near equilibrium  $(0, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 6, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 3, \quad \vec{w}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

## Example 3. Linear dynamics near $(0,0)$


The Linear Approximating System near equilibrium  $(0,0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 6 & 0 \\ 0 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 6, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 3, \quad \vec{w}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Equilibrium  $(0,0)$  is  
a nodal source.

## Example 3. Linear dynamics near $(2, 0)$

The Linear Approximating  
System near equilibrium  $(2, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -6 & -4 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 - 2 \\ x_2 \end{bmatrix}$$

## Example 3. Linear dynamics near $(2, 0)$

The Linear Approximating System near equilibrium  $(2, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -6 & -4 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 - 2 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -6, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 1, \quad \vec{w}_2 = \begin{bmatrix} -4 \\ 7 \end{bmatrix}$$

## Example 3. Linear dynamics near $(2, 0)$


The Linear Approximating System near equilibrium  $(2, 0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -6 & -4 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} x_1 - 2 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -6, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 1, \quad \vec{w}_2 = \begin{bmatrix} -4 \\ 7 \end{bmatrix}$$



Equilibrium  $(2, 0)$  is a saddle

## Example 3. Linear dynamics near $(0, \frac{3}{2})$

The Linear Approximating  
System near equilibrium  $(0, \frac{3}{2})$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -\frac{3}{2} & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - \frac{3}{2} \end{bmatrix}$$

## Example 3. Linear dynamics near $(0, \frac{3}{2})$

The Linear Approximating  
System near equilibrium  $(0, \frac{3}{2})$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -\frac{3}{2} & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - \frac{3}{2} \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 3, \quad \vec{w}_1 = \begin{bmatrix} -4 \\ 1 \end{bmatrix}$$

$$\lambda_2 = -3, \quad \vec{w}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

## Example 3. Linear dynamics near $(0, \frac{3}{2})$


The Linear Approximating System near equilibrium  $(0, \frac{3}{2})$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ -\frac{3}{2} & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - \frac{3}{2} \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 3, \quad \vec{w}_1 = \begin{bmatrix} -4 \\ 1 \end{bmatrix}$$

$$\lambda_2 = -3, \quad \vec{w}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Equilibrium  $(0, \frac{3}{2})$  is a saddle



## Example 3. Linear dynamics near $(\frac{3}{2}, \frac{3}{4})$

The Linear Approximating System  
near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix} \begin{bmatrix} x_1 - \frac{3}{2} \\ x_2 - \frac{3}{4} \end{bmatrix}$$

## Example 3. Linear dynamics near $(\frac{3}{2}, \frac{3}{4})$

The Linear Approximating System  
near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix} \begin{bmatrix} x_1 - \frac{3}{2} \\ x_2 - \frac{3}{4} \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -3 + \frac{3}{2}\sqrt{2} \approx -0.88$$

$$\vec{\mathbf{w}}_1 = \begin{bmatrix} 2 \\ -1 - \sqrt{2} \end{bmatrix}$$

$$\lambda_2 = -3 - \frac{3}{2}\sqrt{2} \approx -5.12$$

$$\vec{\mathbf{w}}_2 = \begin{bmatrix} 2 \\ -1 + \sqrt{2} \end{bmatrix}$$

## Example 3. Linear dynamics near $(\frac{3}{2}, \frac{3}{4})$

The Linear Approximating System  
near equilibrium  $(\frac{3}{2}, \frac{3}{4})$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -\frac{9}{2} & -3 \\ -\frac{3}{4} & -\frac{3}{2} \end{bmatrix} \begin{bmatrix} x_1 - \frac{3}{2} \\ x_2 - \frac{3}{4} \end{bmatrix}$$


Eigenvalues & Eigenvectors:

$$\lambda_1 = -3 + \frac{3}{2}\sqrt{2} \approx -0.88$$

$$\vec{w}_1 = \begin{bmatrix} 2 \\ -1 - \sqrt{2} \end{bmatrix}$$


$$\lambda_2 = -3 - \frac{3}{2}\sqrt{2} \approx -5.12$$

$$\vec{w}_2 = \begin{bmatrix} 2 \\ -1 + \sqrt{2} \end{bmatrix}$$





Equilibrium  $(\frac{3}{2}, \frac{3}{4})$  is  
a nodal sink.


Linear Approx Dynamics Near  $(0, 3/2)$




Linear Approx Dynamics Near  $(3/2, 3/4)$



Linear Approx Dynamics Near  $(0,0)$




Linear Approx Dynamics Near  $(2,0)$



$$\frac{dx_1}{dt} = x_1(6 - 3x_1 - 2x_2)$$

$$\frac{dx_2}{dt} = x_2(3 - x_1 - 2x_2)$$


Nullclines




$$\frac{dx_1}{dt} = x_1(6 - 3x_1 - 2x_2)$$

$$\frac{dx_2}{dt} = x_2(3 - x_1 - 2x_2)$$


Direction Fields on the Nullclines



# Example 3. Global phase portrait




# Example 3. Discussion






# Example 3. Discussion

- The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.




# Example 3. Discussion

- ▶ The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.
- ▶ The co-existence state  $(\frac{3}{2}, \frac{3}{4})$  is asymptotically stable.




### Example 3. Discussion

- ▶ The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.
- ▶ The co-existence state  $(\frac{3}{2}, \frac{3}{4})$  is asymptotically stable.
- ▶ All positive solutions converge to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .




## Example 3. Discussion

- ▶ The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.
- ▶ The co-existence state  $(\frac{3}{2}, \frac{3}{4})$  is asymptotically stable.
- ▶ All positive solutions converge to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .
- ▶ A change of the initial populations does not affect the eventual convergence to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .



## Example 3. Discussion

- ▶ The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.
- ▶ The co-existence state  $(\frac{3}{2}, \frac{3}{4})$  is asymptotically stable.
- ▶ All positive solutions converge to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .
- ▶ A change of the initial populations does not affect the eventual convergence to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .




---

**Question:** Why is the co-existence stable in this system?

## Example 3. Discussion

- ▶ The survival-extinction states  $(2, 0)$  and  $(0, \frac{3}{2})$  are unstable.
- ▶ The co-existence state  $(\frac{3}{2}, \frac{3}{4})$  is asymptotically stable.
- ▶ All positive solutions converge to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .
- ▶ A change of the initial populations does not affect the eventual convergence to the co-existence state  $(\frac{3}{2}, \frac{3}{4})$ .




---

**Question:** Why is the co-existence stable in this system?

**Answer:** Weak competition.

## Example 3. (continued. Weak competition)

$$\begin{cases} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{cases}$$




## Example 3. (continued. Weak competition)

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - x_1 - 2x_2) \end{cases}$$

The competition terms

$-2x_2$  and  $-x_1$





## Example 3. (continued. Weak competition)


$$\begin{cases} x_1' = x_1(6 - 3x_1 - 2x_2) \\ x_2' = x_2(3 - x_1 - 2x_2) \end{cases}$$

The competition terms

$-2x_2$  and  $-x_1$

the resource inhibition terms

$-3x_1$  and  $-2x_2$



## Example 3. (continued. Weak competition)

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - x_1 - 2x_2) \end{cases}$$


The competition terms

$-2x_2$  and  $-x_1$

are “weaker” than

the resource inhibition terms

$-3x_1$  and  $-2x_2$



## Example 3. (continued. Weak competition)

$$\begin{cases} x'_1 = x_1(6 - 3x_1 - 2x_2) \\ x'_2 = x_2(3 - x_1 - 2x_2) \end{cases}$$


The competition terms

$$-2x_2 \text{ and } -x_1$$

are “weaker” than

the resource inhibition terms

$$-3x_1 \text{ and } -2x_2$$



$$\det \begin{bmatrix} 3 & 2 \\ 1 & 2 \end{bmatrix} > 0 \Rightarrow \text{Weak competition} \Rightarrow \text{Stable co-existence}$$

## Example 4. Strong Competition Model.

Competing Species:

$$\begin{cases} x'_1 = x_1(3 - x_1 - 2x_2) \\ x'_2 = x_2(2 - x_1 - x_2) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

## Example 4. Strong Competition Model.

Competing Species:

$$\begin{cases} x_1' = x_1(3 - x_1 - 2x_2) \\ x_2' = x_2(2 - x_1 - x_2) \end{cases}$$

$x_2$  reduces the growth of  $x_1$

$x_1$  reduces the growth of  $x_2$

Equilibria:

$$\begin{cases} x_1(3 - x_1 - 2x_2) = 0 \\ x_2(2 - x_1 - x_2) = 0 \end{cases} \implies \text{Separate to four combinations}$$

$$\begin{cases} x_1 = 0 \\ x_2 = 0 \end{cases}$$

$$\begin{cases} 3 - x_1 - 2x_2 = 0 \\ x_2 = 0 \end{cases}$$

$$\begin{cases} x_1 = 0 \\ 2 - x_1 - x_2 = 0 \end{cases}$$

$$\begin{cases} 3 - x_1 - 2x_2 = 0 \\ 2 - x_1 - x_2 = 0 \end{cases}$$

Four equilibria:  $(x_1, x_2) = (0, 0), (3, 0), (0, 2), (1, 1).$

## Example 4. Linear dynamics near $(0, 0)$

The Linear Approximating  
System near equilibrium  $(0, 0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

## Example 4. Linear dynamics near $(0, 0)$

The Linear Approximating  
System near equilibrium  $(0, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 3, \quad \vec{\mathbf{w}}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 2, \quad \vec{\mathbf{w}}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

## Example 4. Linear dynamics near $(0,0)$


The Linear Approximating System near equilibrium  $(0,0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} 3 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = 3, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = 2, \quad \vec{w}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Equilibrium  $(0,0)$  is  
a nodal source.



## Example 4. Linear dynamics near $(3, 0)$

The Linear Approximating  
System near equilibrium  $(3, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -3 & -6 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 3 \\ x_2 \end{bmatrix}$$

## Example 4. Linear dynamics near $(3, 0)$

The Linear Approximating  
System near equilibrium  $(3, 0)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -3 & -6 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 3 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -3, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = -1, \quad \vec{w}_2 = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$$

## Example 4. Linear dynamics near $(3, 0)$


The Linear Approximating System near equilibrium  $(3, 0)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -3 & -6 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 3 \\ x_2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -3, \quad \vec{w}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$$

$$\lambda_2 = -1, \quad \vec{w}_2 = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$$



Equilibrium  $(3, 0)$  is  
a nodal sink

## Example 4. Linear dynamics near $(0, 2)$

The Linear Approximating  
System near equilibrium  $(0, 2)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - 2 \end{bmatrix}$$

## Example 4. Linear dynamics near $(0, 2)$

The Linear Approximating  
System near equilibrium  $(0, 2)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - 2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -2, \quad \vec{w}_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\lambda_2 = -1, \quad \vec{w}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

## Example 4. Linear dynamics near $(0, 2)$


The Linear Approximating System near equilibrium  $(0, 2)$ :

$$\begin{bmatrix} x_1' \\ x_2' \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ -2 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 - 2 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -2, \quad \vec{w}_1 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$\lambda_2 = -1, \quad \vec{w}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$



Equilibrium  $(0, 2)$  is  
a nodal sink

## Example 4. Linear dynamics near $(1, 1)$

The Linear Approximating System  
near equilibrium  $(1, 1)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix}$$

## Example 4. Linear dynamics near $(1, 1)$

The Linear Approximating System  
near equilibrium  $(1, 1)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix}$$

Eigenvalues & Eigenvectors:

$$\lambda_1 = -1 + \sqrt{2} > 0$$

$$\vec{w}_1 = \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$$

$$\lambda_2 = -1 - \sqrt{2} < 0$$

$$\vec{w}_2 = \begin{bmatrix} \sqrt{2} \\ 1 \end{bmatrix}$$



## Example 4. Linear dynamics near $(1, 1)$

The Linear Approximating System  
near equilibrium  $(1, 1)$ :

$$\begin{bmatrix} x'_1 \\ x'_2 \end{bmatrix} = \begin{bmatrix} -1 & -2 \\ -1 & -1 \end{bmatrix} \begin{bmatrix} x_1 - 1 \\ x_2 - 1 \end{bmatrix}$$


Eigenvalues & Eigenvectors:

$$\lambda_1 = -1 + \sqrt{2} > 0$$

$$\vec{w}_1 = \begin{bmatrix} -\sqrt{2} \\ 1 \end{bmatrix}$$


$$\lambda_2 = -1 - \sqrt{2} < 0$$

$$\vec{w}_2 = \begin{bmatrix} \sqrt{2} \\ 1 \end{bmatrix}$$




Equilibrium  $(1, 1)$  is a saddle



Linear Approx Dynamics Near (0,2)




Linear Approx Dynamics Near (1,1)



Linear Approx Dynamics Near (0,0)




Linear Approx Dynamics Near (3,0)



$$\frac{dx_1}{dt} = x_1(3 - x_1 - 2x_2)$$

$$\frac{dx_2}{dt} = x_2(2 - x_1 - x_2)$$


Nullclines




$$\frac{dx_1}{dt} = x_1(3 - x_1 - 2x_2)$$

$$\frac{dx_2}{dt} = x_2(2 - x_1 - x_2)$$


Direction Fields on the Nullclines



## Example 4. Global phase portrait




# Example 4. Discussion




# Example 4. Discussion

- The survival-extinction states  $(3, 0)$  and  $(0, 2)$  are both asymptotically stable.



# Example 4. Discussion


- ▶ The survival-extinction states  $(3, 0)$  and  $(0, 2)$  are both asymptotically stable.
- ▶ The co-existence state  $(1, 1)$  is unstable.






## Example 4. Discussion

- ▶ The survival-extinction states  $(3, 0)$  and  $(0, 2)$  are both asymptotically stable.
- ▶ The co-existence state  $(1, 1)$  is unstable.
- ▶ Almost all positive solutions converge to either  $(3, 0)$  or  $(0, 2)$ .



## Example 4. Discussion

- ▶ The survival-extinction states  $(3, 0)$  and  $(0, 2)$  are both asymptotically stable.
- ▶ The co-existence state  $(1, 1)$  is unstable.
- ▶ Almost all positive solutions converge to either  $(3, 0)$  or  $(0, 2)$ .
- ▶ A small difference in the initial conditions may make a huge difference in a species' destiny.



## Example 4. (continued. Strong competition)


A small difference in the initial conditions may make a huge difference in a species' destiny.

---

Initial data:

$$x_1(0) = 2.01, x_2(0) = 1.64.$$

As  $t \rightarrow \infty$ ,  $(x_1, x_2) \rightarrow (3, 0)$ .



## Example 4. (continued. Strong competition)


A small difference in the initial conditions may make a huge difference in a species' destiny.

---

Initial data:

$$x_1(0) = 2.01, x_2(0) = 1.64.$$


As  $t \rightarrow \infty$ ,  $(x_1, x_2) \rightarrow (3, 0)$ .



Initial data:


$$x_1(0) = 2.01, x_2(0) = 1.65.$$

As  $t \rightarrow \infty$ ,  $(x_1, x_2) \rightarrow (0, 2)$ .



## Example 4. (continued. Strong competition)


Question: Why is the co-existence unstable in this system?



## Example 4. (continued. Strong competition)

**Question:** Why is the co-existence unstable in this system?

**Answer:** Strong competition.




## Example 4. (continued. Strong competition)

**Question:** Why is the co-existence unstable in this system?

**Answer:** Strong competition.

$$\begin{cases} x_1' = x_1(3 - x_1 - 2x_2) \\ x_2' = x_2(2 - x_1 - x_2) \end{cases}$$



## Example 4. (continued. Strong competition)


**Question:** Why is the co-existence unstable in this system?

**Answer:** Strong competition.

$$\begin{cases} x_1' = x_1(3 - x_1 - 2x_2) \\ x_2' = x_2(2 - x_1 - x_2) \end{cases}$$

The competition terms

$-2x_2$  and  $-x_1$





### Example 4. (continued. Strong competition)

**Question:** Why is the co-existence unstable in this system?


**Answer: Strong competition.**

$$\begin{cases} x'_1 = x_1 \begin{pmatrix} 3 & -x_1 & -2x_2 \end{pmatrix} \\ x'_2 = x_2 \begin{pmatrix} 2 & -x_1 & -x_2 \end{pmatrix} \end{cases}$$

## The competition terms

 $-2x_2$  and  $-x_1$ 

the resource inhibition terms

 $-x_1$  and  $-x_2$ 

## Example 4. (continued. Strong competition)

Question: Why is the co-existence unstable in this system?

Answer: Strong competition.

$$\begin{cases} x_1' = x_1(3 - x_1 - 2x_2) \\ x_2' = x_2(2 - x_1 - x_2) \end{cases}$$


The competition terms

$-2x_2$  and  $-x_1$

**ARE "STRONGER" THAN**

the resource inhibition terms

$-x_1$  and  $-x_2$



## Example 4. (continued. Strong competition)

Question: Why is the co-existence unstable in this system?

Answer: Strong competition.

$$\begin{cases} x'_1 = x_1(3 - x_1 - 2x_2) \\ x'_2 = x_2(2 - x_1 - x_2) \end{cases}$$


The competition terms

$$-2x_2 \text{ and } -x_1$$

**ARE "STRONGER" THAN**

the resource inhibition terms

$$-x_1 \text{ and } -x_2$$



$$\det \begin{bmatrix} 1 & 2 \\ 1 & 1 \end{bmatrix} < 0 \Rightarrow \text{Strong competition} \Rightarrow \begin{cases} \text{One species survives,} \\ \text{the other extincts.} \end{cases}$$